



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS  
 General Certificate of Education  
 Advanced Subsidiary Level and Advanced Level

CANDIDATE  
NAME

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--



**CHEMISTRY**

**9701/34**

Advanced Practical Skills 2

**May/June 2012**

**2 hours**

Candidates answer on the Question Paper.

Additional Materials: As listed in the Confidential Instructions

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Give details of the practical session and laboratory where appropriate, in the boxes provided.

Write in dark blue or black pen.

You may use a soft pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

**DO NOT WRITE IN ANY BARCODES.**

Answer **all** questions.

You may lose marks if you do not show your working or if you do not use appropriate units.

Use of a Data Booklet is unnecessary.

Qualitative Analysis Notes are printed on pages 10 and 11.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

Session
Laboratory

For Examiner's Use	
1	
2	
<b>Total</b>	

This document consists of **11** printed pages and **1** blank page.



1 You are to determine the percentage by mass of sodium ethanedioate in a mixture of sodium ethanedioate and ethanedioic acid.

This experiment involves two steps.

In step one, you will carry out a titration to find the amount of acid,  $C_2O_4H_2$ , present in **FB 3**.  
In step two, you will carry out a second titration to find the total amount of ethanedioate ion,  $C_2O_4^{2-}$ , present in **FB 3**.

Finally, you will use the values found in the two steps to calculate the percentage by mass of sodium ethanedioate in **FB 3**.

**FB 1** is  $0.100\text{ mol dm}^{-3}$  sodium hydroxide, NaOH.

**FB 2** is  $0.0200\text{ mol dm}^{-3}$  potassium manganate(VII),  $KMnO_4$ .

**FB 3** is a mixture of aqueous sodium ethanedioate,  $C_2O_4Na_2$ , and ethanedioic acid,  $C_2O_4H_2$ .

**FB 4** is approximately  $2\text{ mol dm}^{-3}$  sulfuric acid.  
phenolphthalein indicator

**Read through the whole method before starting any practical work.**

**(a) Method**

**Step 1**

- Fill the burette labelled **FB 1** with **FB 1**.
- Pipette  $25.0\text{ cm}^3$  of **FB 3** into a conical flask.
- Add about three drops of phenolphthalein.
- Perform a **rough titration** and record your burette readings in the space below.

The rough titre is .....  $\text{cm}^3$ .

- Carry out as many accurate titrations as you think necessary to obtain consistent results.
- Make certain any recorded results show the precision of your practical work.
- Record in a suitable form below all of your burette readings and the volume of **FB 1** added in each accurate titration.

**Step 2**

- Pipette 25.0 cm<sup>3</sup> of **FB 3** into a conical flask.
- Using a measuring cylinder, add about 25 cm<sup>3</sup> of 2 mol dm<sup>-3</sup> sulfuric acid, **FB 4**, to the flask.
- Place the conical flask on a tripod and gauze and heat to about 80 °C.
- Fill the burette labelled **FB 2** with **FB 2**.
- Use an appropriate method to carefully transfer the hot conical flask onto a white tile under the burette.
- Titrate the mixture in the conical flask with **FB 2** until a permanent pale pink colour is seen. If a permanent brown colour is seen, stop the titration and begin **Step 2** again.
- Perform a **rough titration** and record your burette readings in the space below.

The rough titre is ..... cm<sup>3</sup>.

- Carry out as many accurate titrations as you think necessary to obtain consistent results.
- Make certain any recorded results show the precision of your practical work.
- Record in a suitable form below all of your burette readings and the volume of **FB 2** added in each accurate titration.

I	
II	
III	
IV	
V	
VI	
VII	
VIII	
IX	
X	
XI	
XII	

[12]

**Calculations**

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

**(b) (i)** From your accurate titration results in **Step 1**, obtain a suitable value to be used in your calculations. Show clearly how you have obtained this value.

25.0 cm<sup>3</sup> of **FB 3** required ..... cm<sup>3</sup> of **FB 1**.

(ii) Write an equation for the reaction between sodium hydroxide and ethanedioic acid to give sodium ethanedioate and water.

.....

(iii) Use your answer from (i) to calculate the number of moles of sodium hydroxide, **FB 1**, required to react with 25.0 cm<sup>3</sup> of **FB 3** in **Step 1**.

moles of NaOH = ..... mol

(iv) Use your answer to (iii) to determine the number of moles of ethanedioic acid in 25.0 cm<sup>3</sup> of **FB 3**.

moles of C<sub>2</sub>O<sub>4</sub>H<sub>2</sub> in 25.0 cm<sup>3</sup> of **FB 3** = ..... mol

[2]


(c) (i) From your accurate titration results in **Step 2**, obtain a suitable value to be used in your calculations. Show clearly how you have obtained this value.

25.0 cm<sup>3</sup> of **FB 3** required ..... cm<sup>3</sup> of **FB 2**.

(ii) Use your answer from (i) to calculate the number of moles of potassium manganate(VII), **FB 2**, required to react with 25.0 cm<sup>3</sup> of **FB 3** in **Step 2**.

moles of KMnO<sub>4</sub> = ..... mol

(iii) The equation for the reaction between acidified manganate(VII) ions and ethanedioate ions is shown below.



Calculate the total number of moles of ethanedioate ions in 25.0 cm<sup>3</sup> of **FB 3**.

total moles of C<sub>2</sub>O<sub>4</sub><sup>2-</sup> in 25.0 cm<sup>3</sup> of **FB 3** = ..... mol

(iv) Use your answers to (b)(iv) and (c)(iii) to calculate the number of moles of ethanedioate ions which came from the sodium ethanedioate dissolved in 25.0 cm<sup>3</sup> of **FB 3**.

I	
II	
III	
IV	

moles of C<sub>2</sub>O<sub>4</sub><sup>2-</sup> from C<sub>2</sub>O<sub>4</sub>Na<sub>2</sub> in 25.0 cm<sup>3</sup> of **FB 3** = ..... mol  
[4]

(d) (i) Use your answer to (b)(iv) to calculate the mass of ethanedioic acid, C<sub>2</sub>O<sub>4</sub>H<sub>2</sub>, in 25.0 cm<sup>3</sup> of **FB 3**.

[A<sub>r</sub>: H, 1.0; C, 12.0; O, 16.0]

(If you were unable to answer (b)(iv), you may assume that the number of moles of ethanedioic acid is 6.51 × 10<sup>-4</sup> mol.)

mass of ethanedioic acid = ..... g

(ii) Use your answer to (c)(iv) to calculate the mass of sodium ethanedioate, C<sub>2</sub>O<sub>4</sub>Na<sub>2</sub>, in 25.0 cm<sup>3</sup> of **FB 3**.

[A<sub>r</sub>: C, 12.0; O, 16.0; Na, 23.0]

(If you were unable to answer (c)(iv), you may assume that the number of moles of sodium ethanedioate is 4.13 × 10<sup>-4</sup> mol.)

mass of sodium ethanedioate = ..... g

(iii) Calculate the percentage by mass of sodium ethanedioate present in **FB 3**.

I	
II	
III	
IV	

Percentage by mass of sodium ethanedioate present is ..... %.  
[4]

(e) (i) What is the maximum error in a single burette reading?

maximum error = ..... cm<sup>3</sup>

(ii) A student suggested that using a burette to measure the 25.0 cm<sup>3</sup> of acid would give a more accurate result than using a pipette. The percentage error of a 25.0 cm<sup>3</sup> pipette is 0.24 %. Is the student correct? Explain your answer.

.....  
.....  
.....

[2]

(f) A student decided to use a 25.0 cm<sup>3</sup> pipette instead of a measuring cylinder to measure the volume of **FB 4** in **Step 2**.

State and explain whether this alteration will improve the accuracy of the calculation of the percentage by mass of sodium ethanedioate in the mixture.

.....  
.....  
.....

[1]

[Total: 25]

## 2 Qualitative analysis

At each stage of any test you are to record details of the following.

- colour changes seen
- the formation of any precipitate
- the solubility of such precipitates in an excess of the reagent added

Where gases are released they should be identified by a test, **described in the appropriate place in your observations**.

You should indicate clearly at what stage in a test a change occurs.

Marks are **not** given for chemical equations.

**No additional tests for ions present should be attempted.**

**If any solution is warmed, a boiling tube MUST be used.**

Rinse and reuse test-tubes and boiling tubes where possible.

**Where reagents are selected for use in a test the full name or correct formula of the reagent must be given.**

(a) Compounds **FB 5**, **FB 6** and **FB 7** contain the same non-metal but in three different oxidation states. You are provided with solid samples of **FB 5**, **FB 6** and **FB 7**. Carry out the tests described below and record your observations in the table.

<i>test</i>	<i>observations</i>
(i) To 1 cm depth of dilute hydrochloric acid in a test-tube add a small spatula measure of <b>FB 5</b> .	
(ii) To 1 cm depth of dilute sulfuric acid in a boiling tube add the same depth of aqueous potassium iodide. Add a small spatula measure of <b>FB 5</b> .	
(iii) To 1 cm depth of dilute sulfuric acid in a test-tube add about ten drops of aqueous potassium manganate(VII). Add a small spatula measure of <b>FB 5</b> .	

test	observations
(iv) Place a small spatula measure of <b>FB 6</b> into a hard glass test-tube. Heat the contents gently.	
(v) Place a small spatula measure of <b>FB 6</b> into a boiling tube. Dissolve the solid in 1 cm depth of distilled water. Add 1 cm depth of aqueous sodium hydroxide. Warm the mixture <b>with care</b> .	
(vi) Place a small spatula measure of <b>FB 7</b> into a hard glass test-tube. Heat the contents gently at first, then heat more strongly. Allow to stand for a few minutes	
(vii) Place a small spatula measure of <b>FB 7</b> into a boiling tube. Dissolve the solid in about 1 cm depth of distilled water. Add 1 cm depth of aqueous sodium hydroxide. Warm the mixture <b>with care</b> .	

I	
II	
III	
IV	
V	
VI	

[6]

(b) (i) From your observations in (a), identify the non-metal present in **FB 5**, **FB 6** and **FB 7**.

.....

(ii) Suggest the oxidation state of the non-metal in **FB 5** and **FB 6**.

The oxidation state of the non-metal in **FB 5** is .....

The oxidation state of the non-metal in **FB 6** is .....

(iii) Suggest the type of reaction occurring in (a)(iii).

.....

[3]

(c) Solid compounds containing  $\text{Fe}^{2+}$  and  $\text{Ni}^{2+}$  are usually green. One of these ions is present in **FB 8** and the other in **FB 9**. Both **FB8** and **FB9** are aqueous solutions.

(i) Use the Qualitative Analysis Notes on page 10 to select two reagents that, **used in separate tests**, could identify the presence of the  $\text{Fe}^{2+}$  ion.

The reagents are .....

and .....

(ii) Use your chosen reagents to carry out tests on **both FB 8 and FB 9**. Record your results in an appropriate form in the space below.

(iii) From the results of the tests in (ii), state which solution contains the iron(II) ions.

$\text{Fe}^{2+}$  ions are contained in solution .....

Explain how your observations support your conclusion.

.....  
.....

(iv) Aqueous EDTA is a reagent used to identify some transition metals. To 1 cm depth of the solution containing the nickel(II) ion, add 1 cm depth of aqueous EDTA.

observation

.....  
.....

(v) State what you would expect to **see** if acidified potassium manganate(VII) was added to a sample of the solution containing the iron(II) ion.  
**Do not carry out this experiment.**

expected observation .....

I	
II	
III	
IV	
V	
VI	

[6]

[Total: 15]

## Qualitative Analysis Notes

Key: [ppt. = precipitate]

### 1 Reactions of aqueous cations

ion	reaction with	
	NaOH(aq)	NH <sub>3</sub> (aq)
aluminium, Al <sup>3+</sup> (aq)	white ppt. soluble in excess	white ppt. insoluble in excess
ammonium, NH <sub>4</sub> <sup>+</sup> (aq)	no ppt. ammonia produced on heating	—
barium, Ba <sup>2+</sup> (aq)	no ppt. (if reagents are pure)	no ppt.
calcium, Ca <sup>2+</sup> (aq)	white ppt. with high [Ca <sup>2+</sup> (aq)]	no ppt.
chromium(III), Cr <sup>3+</sup> (aq)	grey-green ppt. soluble in excess giving dark green solution	grey-green ppt. insoluble in excess
copper(II), Cu <sup>2+</sup> (aq)	pale blue ppt. insoluble in excess	blue ppt. soluble in excess giving dark blue solution
iron(II), Fe <sup>2+</sup> (aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess
iron(III), Fe <sup>3+</sup> (aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess
lead(II), Pb <sup>2+</sup> (aq)	white ppt. soluble in excess	white ppt. insoluble in excess
magnesium, Mg <sup>2+</sup> (aq)	white ppt. insoluble in excess	white ppt. insoluble in excess
manganese(II), Mn <sup>2+</sup> (aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess
zinc, Zn <sup>2+</sup> (aq)	white ppt. soluble in excess	white ppt. soluble in excess

[Lead(II) ions can be distinguished from aluminium ions by the insolubility of lead(II) chloride.]

## 2 Reactions of anions

ion	reaction
carbonate, $\text{CO}_3^{2-}$	$\text{CO}_2$ liberated by dilute acids
chromate(VI), $\text{CrO}_4^{2-}(\text{aq})$	yellow solution turns orange with $\text{H}^+(\text{aq})$ ; gives yellow ppt. with $\text{Ba}^{2+}(\text{aq})$ ; gives bright yellow ppt. with $\text{Pb}^{2+}(\text{aq})$
chloride, $\text{Cl}^-(\text{aq})$	gives white ppt. with $\text{Ag}^+(\text{aq})$ (soluble in $\text{NH}_3(\text{aq})$ ); gives white ppt. with $\text{Pb}^{2+}(\text{aq})$
bromide, $\text{Br}^-(\text{aq})$	gives cream ppt. with $\text{Ag}^+(\text{aq})$ (partially soluble in $\text{NH}_3(\text{aq})$ ); gives white ppt. with $\text{Pb}^{2+}(\text{aq})$
iodide, $\text{I}^-(\text{aq})$	gives yellow ppt. with $\text{Ag}^+(\text{aq})$ (insoluble in $\text{NH}_3(\text{aq})$ ); gives yellow ppt. with $\text{Pb}^{2+}(\text{aq})$
nitrate, $\text{NO}_3^-(\text{aq})$	$\text{NH}_3$ liberated on heating with $\text{OH}^-(\text{aq})$ and $\text{Al}$ foil
nitrite, $\text{NO}_2^-(\text{aq})$	$\text{NH}_3$ liberated on heating with $\text{OH}^-(\text{aq})$ and $\text{Al}$ foil; NO liberated by dilute acids (colourless NO $\rightarrow$ (pale) brown $\text{NO}_2$ in air)
sulfate, $\text{SO}_4^{2-}(\text{aq})$	gives white ppt. with $\text{Ba}^{2+}(\text{aq})$ or with $\text{Pb}^{2+}(\text{aq})$ (insoluble in excess dilute strong acids)
sulfite, $\text{SO}_3^{2-}(\text{aq})$	$\text{SO}_2$ liberated with dilute acids; gives white ppt. with $\text{Ba}^{2+}(\text{aq})$ (soluble in excess dilute strong acids)

## 3 Tests for gases

gas	test and test result
ammonia, $\text{NH}_3$	turns damp red litmus paper blue
carbon dioxide, $\text{CO}_2$	gives a white ppt. with limewater (ppt. dissolves with excess $\text{CO}_2$ )
chlorine, $\text{Cl}_2$	bleaches damp litmus paper
hydrogen, $\text{H}_2$	“pops” with a lighted splint
oxygen, $\text{O}_2$	relights a glowing splint
sulfur dioxide, $\text{SO}_2$	turns acidified aqueous potassium dichromate(VI) from orange to green

---

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.