



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS  
General Certificate of Education  
Advanced Subsidiary Level and Advanced Level

CANDIDATE  
NAME

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--

**CHEMISTRY**

**9701/21**

Paper 2 Structured Questions AS Core

**May/June 2013**

**1 hour 15 minutes**

Candidates answer on the Question Paper.

Additional Materials: Data Booklet

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a soft pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

**DO NOT WRITE IN ANY BARCODES.**

Answer **all** questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

A Data Booklet is provided.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

For Examiner's Use	
1	
2	
3	
4	
5	
Total	

This document consists of **10** printed pages and **2** blank pages.



Answer **all** the questions in the spaces provided.

For  
Examiner's  
Use

- 1** A sample of a fertiliser was known to contain ammonium sulfate,  $(\text{NH}_4)_2\text{SO}_4$ , and sand only.

A 2.96 g sample of the solid fertiliser was heated with  $40.0 \text{ cm}^3$  of  $\text{NaOH}(\text{aq})$ , an excess, and all of the ammonia produced was boiled away.

After cooling, the remaining  $\text{NaOH}(\text{aq})$  was exactly neutralised by  $29.5 \text{ cm}^3$  of  $2.00 \text{ mol dm}^{-3}$   $\text{HCl}$ .

In a separate experiment,  $40.0 \text{ cm}^3$  of the original  $\text{NaOH}(\text{aq})$  was exactly neutralised by  $39.2 \text{ cm}^3$  of the  $2.00 \text{ mol dm}^{-3}$   $\text{HCl}$ .

- (a) (i)** Write balanced equations for the following reactions.

$\text{NaOH}$  with  $\text{HCl}$

.....

$(\text{NH}_4)_2\text{SO}_4$  with  $\text{NaOH}$

.....

- (ii)** Calculate the amount, in moles, of  $\text{NaOH}$  present in the  $40.0 \text{ cm}^3$  of the original  $\text{NaOH}(\text{aq})$  that was neutralised by  $39.2 \text{ cm}^3$  of  $2.00 \text{ mol dm}^{-3}$   $\text{HCl}$ .

- (iii)** Calculate the amount, in moles, of  $\text{NaOH}$  present in the  $40.0 \text{ cm}^3$  of  $\text{NaOH}(\text{aq})$  that remained after boiling the  $(\text{NH}_4)_2\text{SO}_4$ .

- (iv)** Use your answers to **(ii)** and **(iii)** to calculate the amount, in moles, of  $\text{NaOH}$  that reacted with the  $(\text{NH}_4)_2\text{SO}_4$ .

- (v) Use your answers to (i) and (iv) to calculate the amount, in moles, of  $(\text{NH}_4)_2\text{SO}_4$  that reacted with the NaOH.

- (vi) Hence calculate the mass of  $(\text{NH}_4)_2\text{SO}_4$  that reacted.

- (vii) Use your answer to (vi) to calculate the percentage, by mass, of  $(\text{NH}_4)_2\text{SO}_4$  present in the fertiliser.  
Write your answer to a suitable number of significant figures.

[9]

- (b) The uncontrolled use of nitrogenous fertilisers can cause environmental damage to lakes and streams. This is known as *eutrophication*.

What are the processes that occur when excessive amounts of nitrogenous fertilisers get into lakes and streams?


.....  
 .....  
 ..... [2]

- (c) Large quantities of ammonia are manufactured by the Haber process.  
Not all of this ammonia is used to make fertilisers.  
State **one** large-scale use for ammonia, **other than** in the production of nitrogenous fertilisers.

..... [1]

[Total: 12]

- 2 Ammonium nitrate fertiliser is manufactured from ammonia. The first reaction in the manufacture of the fertiliser is the catalytic oxidation of ammonia to form nitrogen monoxide, NO. This is carried out at about  $1 \times 10^3$  kPa (10 atmospheres) pressure and a temperature of 700 to 850 °C.



- (a) Write the expression for the equilibrium constant,  $K_p$ , stating the units.

$K_p =$

units .....

[2]

- (b) What will be the effect on the yield of NO of **each** of the following?  
In each case, explain your answer.

- (i) increasing the temperature

.....  
.....  
.....

- (ii) decreasing the applied pressure

.....  
.....  
.....

[4]

(c) The standard enthalpy changes of formation of  $\text{NH}_3(\text{g})$  and  $\text{H}_2\text{O}(\text{g})$  are as follows.

$$\text{NH}_3(\text{g}), \Delta H_f^\circ = -46.0 \text{ kJ mol}^{-1}$$

$$\text{H}_2\text{O}(\text{g}), \Delta H_f^\circ = -242 \text{ kJ mol}^{-1}$$

Use these data and the value of  $\Delta H_{\text{reaction}}^\circ$  given below to calculate the standard enthalpy change of formation of  $\text{NO}(\text{g})$ .

Include a sign in your answer.



[4]

[Total: 10]

- 3 This question refers to the elements in the section of the Periodic Table shown below.

For  
Examiner's  
Use

		H						He	
Li	Be			B	C	N	O	F	Ne
Na	Mg			Al	Si	P	S	Cl	Ar
K	Ca	..... transition elements .....		Ga	Ge	As	Se	Br	Kr

- (a) From this list of elements, identify in **each** case **one** element that has the property described. Give the **symbol** of the element.

- (i) An element that floats on cold water and reacts readily with it.

.....

- (ii) An element that forms an oxide that is a reducing agent.

.....

- (iii) The element that has the smallest first ionisation energy.

.....

- (iv) The element which has a giant molecular structure **and** forms an oxide which has a simple molecular structure.

.....

- (v) The element in Period 3 (Na to Ar) that has the smallest anion.

.....

- (vi) The element in Period 3 (Na to Ar) which forms a chloride with a low melting point and an oxide with a very high melting point.

.....

[6]

- (b) Use the elements in Period 3 (Na to Ar) in the section of the Periodic Table opposite to identify the oxide(s) referred to below.  
In **each** case, give the **formula** of the oxide(s).

- (i) An oxide which when placed in water for a long time has no reaction with it.

.....

- (ii) An oxide which dissolves readily in water to give a strongly alkaline solution.

.....

- (iii) Two acidic oxides formed by the same element.

..... and .....

- (iv) An oxide which is amphoteric.

.....

[5]

- (c) Fluorine reacts with other elements in Group VII to form a number of different compounds.  
Two such compounds and their boiling points are given in the table.

compound	$\text{ClF}_3$	$\text{BrF}_3$
boiling point/ $^{\circ}\text{C}$	12	127

- (i) The two molecules have similar electronic configurations.  
Showing outer electrons only, draw a 'dot-and-cross' diagram of the bonding in  $\text{ClF}_3$ .

- (ii) The two molecules have the same shape.  
Suggest why the boiling points are significantly different.

.....  
.....  
.....  
.....

[4]

[Total: 15]





- 4 Organic chemistry is the chemistry of carbon compounds. The types of organic reactions that you have studied are listed below.

For  
Examiner's  
Use

addition

elimination

hydrolysis

oxidation

reduction

substitution

Addition and substitution reactions are further described as follows.

electrophilic

nucleophilic

free radical

Complete the table below.

Fill in the central column by using **only** the types of reaction given in the lists above.

Use **both** lists when appropriate.

In the right hand column give the formula(e) of the reagent(s) you would use to carry out the reaction given.

organic reaction	type of reaction	reagent(s)
$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{Br} \rightarrow$ $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2$		
$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH} \rightarrow$ $\text{BrCH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$		
$\text{CH}_3\text{COCH}_3 \rightarrow$ $\text{CH}_3\text{C}(\text{OH})(\text{CN})\text{CH}_3$		
$\text{CH}_3\text{CH}(\text{OH})\text{CH}_2\text{CH}_3 \rightarrow$ $\text{CH}_3\text{CH}=\text{CHCH}_3$		

[Total: 11]

5 Crotonaldehyde,  $\text{CH}_3\text{CH}=\text{CHCHO}$ , occurs in soybean oils.

- (a) In the boxes below, write the **structural formula** of the organic compound formed when crotonaldehyde is reacted separately with each reagent under suitable conditions. If you think no reaction occurs, write 'NO REACTION' in the box.

reaction	reagent	product
A	$\text{Br}_2$ in an inert organic solvent	
B	$\text{PCl}_3$	
C	$\text{H}_2$ and Ni catalyst	
D	$\text{NaBH}_4$	
E	$\text{K}_2\text{Cr}_2\text{O}_7/\text{H}^+$	

[5]

- (b) Crotonaldehyde exists in more than one stereoisomeric form. Draw the **displayed formulae** of the **stereoisomers** of crotonaldehyde. Label **each** isomer.

[3]

(c) Draw the **skeletal formula** of crotonaldehyde.

[1]

(d) The product of reaction E in the table opposite will react with a solution containing acidified manganate(VII) ions.

Draw the **structural formulae** of the organic products when the reagent is

(i) cold, dilute;

(ii) hot, concentrated.

[3]

[Total: 12]

---

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.